Jessica Barrow BSc Physiotherapy (WITS) 083 256 0434



Room GF03 Waterfall Hospital Cnr. Magwa Crescent and Mac Mac Avenue Tel: 011 304-7829 Fax: 011 304-7941

DR. MICHAEL BARROW PCL AND POSTEROLATERAL CORNER RECONSTRUCTION – REHABILITATION PROTOCOL

	POST OP WEEK				POST OP MONTH		
	1 to 2	3 to 4	5 to 6	7 to 12	4 to 5	6 to 9	9 to 12
Brace	In situ	In situ	In situ	-	-	-	-
Weight-bearing (normal gait pattern ASAP)	See post-op instructions			FWB	FWB	FWB	FWB
ROM Goal	0° to 70°	0° to 70°	0° to 70°	0° to Full	0° to Full	0° to Full	0° to Full
ROM Exercises (within set ROM)							
Active assisted knee flex (foot sliding)	-	-	-	0° to 110°*			
Active knee extension	v	٧	v	v			
Passive knee flex (proximal tibial force directed anteriorly)	v	٧	v	v			
Oedema management (RICE)	V	V	V	As reqd.	As reqd.	As reqd.	As reqd.

	POST OP WEEK				POST OP MONTH			
	1 to 2	3 to 4	5 to 6	7 to 12	4 to 5	6 to 9	9 to 12	
Stretching Hamstrings, calf, ITB,	٧	v	V	V	V	v	v	
Patella mobilisations	٧	٧	V	v	As reqd.	As reqd.	As reqd.	
Strengthening Isometric quads / SLR	V	V	V	√	V	V	√	
Closed chain (gait re-ed, mini squats within ROM limits, toe standing)			V	v	٧	v	v	
Wall slides				0° to 60°	v	v	v	
Open chain knee extension				25° to 0°	25° to 0°	70° to 0°	90° to 0°	
Hamstring curls						v	v	
Leg press				v	v	v	v	
Hip extension, ABD, ADD (avoid stresses on knee)	v	٧	v	v	v	v	v	
Cardiovascular stationary bicycle (when 100° flexion)				v	V	V	٧	
Step machine / swimming – straight kick / elliptical machine					v	v	v	
Running – straight						v	v	
Proprioception (e.g. weight transfers, balance board, mini tramp)				v	٧	V	٧	
Dynamic stability (e.g. stepping on / off different surfaces and in all directions)				v	v	v	v	

	POST OP WEEK				POST OP MONTH		
	1 to 2	3 to 4	5 to 6	7 to 12	4 to 5	6 to 9	9 to 12
Sport specific / agility drills (e.g. shuttle runs, cariocas, figure 8's)						V **	V
Plyometrics (e.g. bounding, hopping, jumping)						V**	v
Activities							
Work – sedentary		٧	v	v	٧	٧	v
Work – heavy				V***	V***	٧	v
Driving				8 / 52	٧	٧	٧
Full sports							√ **

*Active knee flex may only begin at 6 weeks.

**Progressed within limits of pain, swelling and muscle control.

***Dependent on type of employment / sport